arXiv:1012.5612v1 [math.AP] 27 Dec 2010

Solutions for a Nonhomogeneous Nonlinear Schroedinger Equation with Double Power Nonlinearity

M.Ghimenti [*], A.M.Micheletti [†]

Abstract

We consider the problem $-\Delta u + V(x)u = f'(u) + g(x)$ in $\mathbb{R}^N$, under the assumption $\lim_{x\to\infty} V(x) = 0$, and with the non linear term f with a *double power* behavior. We prove the existence two solutions when g is sufficiently small and $V < 0$.

Keywords: Nonlinear Equations, Variational Methods, Orlicz Spaces

1 Perturbation of NSE

We consider the existence of solutions of the following nonhomogeneous problem

$$\begin{cases} -\Delta u + V(x)u = f'(u) + g(x), & x \in \mathbb{R}^N; \\ E_g^V(u) < \infty. \end{cases} \tag{$\mathscr{P}$}$$

where the energy functional is defined by

$$E_g(u) = E_g^V(u) = \frac{1}{2}\int_{\mathbb{R}^N} |\nabla u|^2 + V(x)u^2(x)dx - \int_{\mathbb{R}^N} f(u)dx - \int_{\mathbb{R}^N} g(x)u(x)dx.$$

The nonlinearity is given by a function f of *double power* type that is an even function $f \in C^3(\mathbb{R},\mathbb{R})$ with $f(0) = f'(0) = f''(0) = 0$ satisfying the following requirements:

---

[*] Dipartimento di Matematica Applicata, Università di Pisa, via Buonarroti 1c, 56127, Pisa, Italy

[†] Dipartimento di Matematica Applicata, Università di Pisa, via Buonarroti 1c, 56127, Pisa, Italy

1. there exist positive numbers c_0, c_2, p, q with $2 < p < 2^* < q$ such that

$$\begin{cases} c_0|s|^p \leq f(s) & \text{for } |s| \geq 1; \\ c_0|s|^q < f(s) & \text{for } |s| \leq 1; \end{cases} \tag{f_0}$$

$$\begin{cases} |f''(s)| \leq c_2|s|^{p-2} & \text{for } |s| \geq 1; \\ |f''(s)| \leq c_2|s|^{q-2} & \text{for } |s| \leq 1; \end{cases} \tag{f_2}$$

2. there exists $\mu_1 > 2$ and $\mu_2 > 1$ such that, for all $s \neq 0$

$$0 < \mu_1 f(s) \leq f'(s)s, \qquad \mu_2 f'(s)s < f''(s)s^2, \qquad f'''(s)s^3 > 0; \tag{f_μ}$$

3. for any $u \in \mathcal{D}^{1,2}$ we have

$$f'''(u)u^3 \in L^1. \tag{f_3}$$

For example the required assumptions are satisfied by $f(s) = \frac{|s|^q}{1+|s|^{q-p}}$ with $q - p$ small enough, as shown in the appendix.

We assume $V \in L^{N/2}(\mathbb{R}^N) \cap L^t$, for some $t > N/2$ and

$$||V||_{L^{N/2}} < S := \inf_{u \in \mathcal{D}^{1,2}} \frac{\displaystyle\int_{\mathbb{R}^N} |\nabla u|^2}{\left(\displaystyle\int_{\mathbb{R}^N} |u|^{2^*}\right)^{2/2^*}}. \tag{1}$$

Moreover, we want $V \leq 0$ and $V < 0$ on a set of positive measure.

In [18] the existence of two positive solutions $u_1, u_2 \in H^1(\mathbb{R}^N)$ of the equation $-\Delta u + u = |u|^{p-2}u + g$ is proved when $g \in L^2$ satisfies $0 \leq g \leq C\exp(-(1+\varepsilon)|x|)$, $g \not\equiv 0$.

Recently, in [17], a similar problem for the p-laplacian is studied. Namely, the author proves, with variational techniques, that the problem $-\Delta_p u + c|u|^{p-2}u = |u|^{p^*-2}u + f(x,u) + h(x)$ in $\mathbb{R}^N$, where $2 \leq p < N$, $c > 0$, $h \in W^{-1,p'}(\mathbb{R}^N)$ and f a is continuous superlinear function such that $f(x,0) = 0$ and $f(x,u) = o(|u|^{p^*-1})$ as $|u| \to \infty$, admits two positive solutions $u_1, u_2 \in H^1(\mathbb{R}^N)$.

The existence of a positive solution of the problem $-\Delta u + u = |u|^{p-1}u + g$ on $\mathbb{R}^N$, $u(x) \to 0$ for $|x| \to \infty$, was proven in [10] when $p > \frac{N}{N-2}$ and $g \in C^{0,\alpha}(\mathbb{R}^N)$, $g \geq 0$, $g \not\equiv 0$ and $g(x) \leq \frac{C}{(1+|x|^2)^{p/p-1}}$ for some $C > 0$. In [3] there is a result of multiplicity for this problem.

In [16] the author shows that the Dirichlet problem on a bounded domain $\Omega \subset \mathbb{R}^N$ in the critical case $-\Delta u = |u|^{2^*-2}u + g$ has two solutions $u_0, u_1 \in$

$H_0^1(\Omega)$, for g satisfying a suitable condition, and if $g \geq 0$ then $u_0 \geq 0$ and $u_1 \geq 0$.

We are interested in studying the problem with double power nonlinearity.

In pioneering work Berestycki and Lions [7, 8] showed the existence of a positive solution in the case $V \equiv 0$ when $f''(0) = 0$, f has a supercritical growth near the origin and subcritical at infinity.

More recently in the papers [2, 5, 6, 14] the double-power growth condition has been used to obtain the existence of positive solutions for different problems of the tipe $(\mathscr{P})$. In particular, in [5], the authors proved that in the same hypothesis on V the homogeneous problem

$$-\Delta u + Vu = f'(u) \tag{2}$$

has a ground state solution (i.e. least energy nontrivial solution). Other results on similar problems with the double power nonlinearity can be found in [1, 2, 12].

In this paper we prove the following theorem

Theorem 1. *If $g \in L^{\frac{2N}{N+2}} \cap L^s$, for some $s > \frac{2N}{N+2}$, and if $||g||_{\frac{2N}{N+2}}$ is sufficiently small there exist two solutions of problem $(\mathscr{P})$ in $\mathcal{D}^{1,2}$. The first solution is close to 0; if also $||g||_{L^{p'}\cap L^{q'}}$ is small enough, the critical value of the second solution is close to the least energy level m_V of the homogeneous problem (2).*

Furthermore, if $g \geq 0$ the two solutions are non negative.

Remark 2. Indeed the hypothesis on the sign of V is used only to find the second solution, but we prefer a more compact claim for the theorem. Anyway, in the proofs we focus out when we use any hypothesis.

To get the solutions of $(\mathscr{P})$ we look for critical points of the functional E_g^V constrained on the Nehari manifold

$$\begin{aligned} \mathcal{N}_g^V &= \mathcal{N}_g = \left\{ u \in \mathcal{D}^{1,2} \ : \ \langle \nabla E_g(u), u \rangle = 0, u \neq 0 \right\} = \\ &= \left\{ u \in \mathcal{D}^{1,2} \smallsetminus 0 : \int_{\mathbb{R}^N} |\nabla u|^2 + \int_{\mathbb{R}^N} Vu^2 - \int_{\mathbb{R}^N} f'(u)u - \int_{\mathbb{R}^N} gu = 0 \right\}. \end{aligned}$$

The study of the structure of the Nehari manifold will be a fundamental part of this paper.

This paper is organized as follows: in section 2, we recall some technical results concerning the appropriate function space required by the growth properties of the nonlinearity f . Moreover, we study the geometry and the properties of the Nehari manifold. In section 3, we prove a Splitting Lemma necessary to overcome the lack of compactness. This lemma is a variant for a well known result of [15]. In section 4 we prove the existence of two distinct critical points of the functional on the Nehari manifold.

2 Notations and preliminary result

We will use the following notations

- $\mathcal{D}^{1,2} = \mathcal{D}^{1,2}(\mathbb{R}^N)$ = completion of $C_0^\infty(\mathbb{R}^N)$ with respect to the norm
$$||u|| = \left(\int_{\mathbb{R}^N} |\nabla u|^2 \right)^{1/2};$$
- $||u||_V^2 = \displaystyle\int_{\mathbb{R}^N} |\nabla u|^2 + \int_{\mathbb{R}^N} Vu^2$; notice that, by (1), we have that $||u||_V$ is a norm in $\mathcal{D}^{1,2}$ equivalent to the usual one;
- $2^* = \frac{2N}{N-2}$;
- $m_g = \inf\limits_{u \in \mathcal{N}_g^V} E_g^V(u)$;
- $m_{1,g} = \inf\limits_{u \in \mathcal{N}_g^-} E_g^V(u)$;
- $m_0 = \inf\limits_{u \in \mathcal{N}_0^0} E_0^0(u)$; we call ω the minimizer of E_0^0 on $\mathcal{N}_0^0$ radially symmetric;
- $m_V = \inf\limits_{u \in \mathcal{N}_0^V} E_0^V(u)$; we call $\bar{u}$ the minimizer of E_0^V on $\mathcal{N}_0^V$;
- $\Gamma_u = \{x \in \mathbb{R}^N \ : \ |u(x)| > 1\}$;
- $|A|$ = the Lebesgue measure of the subset $A \subset \mathbb{R}^N$;
- $B_R = \{x \in \mathbb{R}^N \ : \ |x| \leq R\}$;
- $B_R^C = \mathbb{R}^N \setminus B_R$;
- $u_y(x) = u(x+y)$.

In order to study the properties of the functional E_g^V and its Nehari manifold, we consider some suitable Orlicz space $L^p + L^q$, where $2 < p < 2^* < q$, related to the double power growth behavior of the function f. We recall some properties of these spaces to get the smoothness of the functional E_g^V

Given $p \neq q$, we consider the space $L^p + L^q$ made up of the functions $v : \mathbb{R}^N \to \mathbb{R}$ such that

$$v = v_1 + v_2 \quad \text{with} \quad v_1 \in L^p, v_2 \in L^q. \tag{3}$$

The space $L^p + L^q$ is a Banach space equipped with the norm:

$$||v||_{L^p+L^q} = \inf\{\ ||v_1||_{L^p} + ||v_2||_{L^q}\ :\ v_1 \in L^p, v_2 \in L^q,\ v_1 + v_2 = v\}. \tag{4}$$

It is well known (see, for example [9]) that $L^p + L^q$ coincides with the dual of $L^{p'} \cap L^{q'}$. Then:

$$L^p + L^q = \left(L^{p'} \cap L^{q'}\right)' \quad \text{with}\ \ p' = \frac{p}{p-1},\ q' = \frac{q}{q-1}, \tag{5}$$

and we can introduce the following norm equivalent to the previous one

$$||v||_{L^p+L^q} = \inf_{\varphi \neq 0} \frac{\int v\varphi}{||\varphi||_{L^{p'}} + ||\varphi||_{L^{q'}}}. \tag{6}$$

Hereafter we recall some results useful for this paper contained in [4, 6].

Lemma 3. *We have*

1. *if $v \in L^p + L^q$, the following inequalities hold:*

$$\begin{aligned} &\max\left[||v||_{L^q(\mathbb{R}^N\smallsetminus\Gamma_v)} - 1, \frac{1}{1+|\Gamma_v|^{\frac{1}{\tau}}}||v||_{L^p(\Gamma_v)}\right] \leq \\ \leq\ & ||v||_{L^p+L^q} \leq \\ \leq\ & \max[||v||_{L^q(\mathbb{R}^N\smallsetminus\Gamma_v)}, ||v||_{L^p(\Gamma_v)}] \end{aligned}$$

   when $\tau = \frac{pq}{q-p}$;

2. *let $\{v_n\} \subset L^p + L^q$. Then $\{v_n\}$ is bounded in $L^p + L^q$ if and only if the sequences $\{|\Gamma_{v_n}|\}$ and $\{||v||_{L^q(\mathbb{R}^N\smallsetminus\Gamma_{v_n})} + ||v||_{L^p(\Gamma_{v_n})}\}$ are bounded.*

3. *f' is a bounded map from $L^p + L^q$ into $L^{\frac{p}{p-1}} \cap L^{\frac{q}{q-1}}$*

Remark 4. By the previous lemma we have $L^{2^*} \subset L^p + L^q$ when $2 < p < 2^* < q$. Then, by Sobolev inequality, we get the continuous embedding

$$\mathcal{D}^{1,2}(\mathbb{R}^N) \subset L^p + L^q.$$

In order to prove the C^2 regularity of the functional E_g^V, we need the following lemmas proved in [6]

Lemma 5. *If f satisfies the hypothesis (f_0) and (f_2), we have that*

1. *if θ, u are bounded in $L^p + L^q$, then $f''(\theta)u$ is bounded in $L^{p'} \cap L^{q'}$;*

2. *f'' is a bounded map from L^p+L^q into $L^{p/p-2}\cap L^{q/q-2}$;*

3. *f'' is a continuous map from L^p+L^q into $L^{p/p-2}\cap L^{q/q-2}$;*

4. *the map $(u,v)\mapsto uv$ from $(L^p+L^q)^2$ in $L^{p/2}+L^{q/2}$ is bounded.*

Lemma 6. *The functional E_g^V is of class C^2 and it holds*

$$\begin{aligned} E_g'(u)[v]=\langle\nabla E_g^V(u),v\rangle &= \int_{\mathbb{R}^N}\nabla u\nabla v+Vuv-f'(u)v-gv; \qquad (7)\\ E_g''(u)[v,w] &= \int_{\mathbb{R}^N}\nabla v\nabla w+Vvw-f''(u)vw. \qquad (8)\end{aligned}$$

Moreover the Nehari manifold defined as

$$\mathcal{N}_g^V=\left\{u\in\mathcal{D}^{1,2}\smallsetminus 0\ :\ \int_{\mathbb{R}^N}|\nabla u|^2+Vu^2-f'(u)udx-gu=0\right\} \qquad (9)$$

is of class C^1 and its tangent space at the point u is

$$T_u\mathcal{N}_g^V=\left\{v\in\mathcal{D}^{1,2}:\int_{\mathbb{R}^N}2\nabla u\nabla v+2Vuv-f'(u)vdx-f''(u)uv-gv=0\right\}.$$

At last, we introduce the functions

$$\begin{aligned} \varphi_0^u(t) &= \varphi_0(t):=E_0^V(tu)=\int_{\mathbb{R}^N}\frac{t^2}{2}(|\nabla u|^2+Vu^2)-f(tu); \qquad (10)\\ \varphi_g^u(t) &= \varphi_g(t):=E_g^V(tu)=\varphi_0(t)-t\int_{\mathbb{R}^N}gu. \qquad (11)\end{aligned}$$

We have that

$$\begin{aligned} \varphi_g'(t) &= t||u||_V^2-\int_{\mathbb{R}^N}f'(tu)u-\int_{\mathbb{R}^N}gu; \qquad (12)\\ \varphi_g''(t) &= ||u||_V^2-\int_{\mathbb{R}^N}f''(tu)u^2; \qquad (13)\\ \varphi_g'''(t) &= -\int_{\mathbb{R}^N}f'''(tu)u^3. \qquad (14)\end{aligned}$$

Notice that the conditions on f assure that also $\varphi_g'''(t)$ exists. Furthermore, if $\frac{d}{dt}\varphi_g(\bar t)=0$, then $\langle\nabla E(\bar t u),u\rangle=0$, so $\bar t u\in\mathcal{N}_g^V$, and *vice versa*, so we want to find the critical points of $\varphi_g(t)$.

To study the manifold $\mathcal{N}_g^V$ it is useful to consider the following manifold:

$$\mathcal{V}=\left\{w\neq 0: G(w):=||w||_V^2-\int_{\mathbb{R}^N}f''(w)w^2=0\right\}. \qquad (15)$$

Lemma 7. *We have that for all $u \in \mathcal{D}^{1,2}$ there exists an unique $T_u > 0$ such that $T_u u \in \mathcal{V}$*

Proof. We have that, using (f_μ) and (f_0),

$$\begin{aligned}
\varphi_0'(t) &= t||u||_V^2 - \int_{\mathbb{R}^N} f'(tu)u \leq t||u||_V^2 - \frac{\mu_1}{t}\int_{\mathbb{R}^N} f(tu) \leq \\
&\leq t||u||_V^2 - t^{q-1}c_0\mu_1 \int_{t|u|<1} |u|^q - t^{p-1}c_0\mu_1 \int_{t|u|\geq 1} |u|^p \leq \qquad (16)\\
&\leq t||u||_V^2 - t^{p-1}c_0\mu_1 \int_{|u|\geq 1} |u|^p \to -\infty \text{ when } t \to \infty,
\end{aligned}$$

because $p > 2$. Furthermore we have that $\varphi_0'(t)$ is strictly concave when $t \neq 0$, and that $\varphi_0''(0) > 0$, so, for every $u \in \mathcal{D}^{1,2}$ there exist an unique maximum point $T_u > 0$ for the function $\varphi_0'(t)$. Thus

$$0 = T_u^2 \varphi_0''(T_u) = ||T_u u||_V^2 - \int_{\mathbb{R}^N} f''(T_u u)(T_u u)^2.$$

□

Proposition 8. *We have that* $\inf\limits_{w \in \mathcal{V}} ||w||_V^2 > 0$.

Proof. By contradiction, we suppose that there exists a sequence $\{w_n\}_n \subset \mathcal{V}$ such that $||w_n||_V^2$ converges to 0. We set $t_n = ||w_n||_V$, hence we can write $w_n = t_n v_n$ where $||v_n||_V = 1$. Remark 4, the sequence is bounded in $L^p + L^q$. Since $w_n \in \mathcal{V}$ and t_n converges to 0, we have

$$\begin{aligned}
1 &= ||v_n||_V^2 = \frac{||w_n||_V^2}{t_n^2} = \frac{1}{t_n^2}\int\limits_{\mathbb{R}^N} f''(t_n v_n) v_n^2 \leq \\
&\leq c_2 t_n^{q-2} \int\limits_{\mathbb{R}^N \smallsetminus \Gamma_{t_n v_n}} |v_n|^q + c_2 t_n^{p-2} \int\limits_{\Gamma_{t_n v_n}} |v_n|^p \leq \\
&\leq c_2 t_n^{q-2} \int\limits_{\mathbb{R}^N \smallsetminus \Gamma_{t_n v_n}} |v_n|^q + c_2 t_n^{p-2} \int\limits_{\Gamma_{v_n}} |v_n|^p \leq \\
&\leq c_2 t_n^{q-2} \int\limits_{\mathbb{R}^N \smallsetminus \Gamma_{v_n}} |v_n|^q + c_2 t_n^{q-2} \int\limits_{(\mathbb{R}^N \smallsetminus \Gamma_{t_n v_n}) \cap \Gamma_{v_n}} \frac{|v_n|^p}{t_n^{q-p}} + c_2 t_n^{p-2} \int\limits_{\Gamma_{v_n}} |v_n|^p \leq \\
&\leq c_2 t_n^{q-2} \int\limits_{\mathbb{R}^N \smallsetminus \Gamma_{v_n}} |v_n|^q + 2c_2 t_n^{p-2} \int\limits_{\Gamma_{v_n}} |v_n|^p.
\end{aligned}$$

Hence we get

$$1 \leq c_2 t_n^{q-2} \int\limits_{\mathbb{R}^N \setminus \Gamma_{v_n}} |v_n|^q + 2c_2 t_n^{p-2} \int\limits_{\Gamma_{v_n}} |v_n|^p$$

and by claim 2 of Remark 3 we get the contradiction. □

Lemma 9. *Let $u \in \mathcal{D}^{1,2}$ and let T_u the unique positive number such that $T_u u \in \mathcal{V}$. Then*

$$L = \inf_{||u||_V = 1} T_u - \int_{\mathbb{R}^N} f'(T_u u)u > 0. \tag{17}$$

Proof. By contradiction, suppose that there exists a minimizing sequence u_n, with $||u_n||_V = 1$ such that $T_{u_n} - \int f'(T_{u_n} u_n)u_n := \sigma_n \to 0$. Let $w_n = T_{u_n} u_n$. We have that

$$T_{u_n}^2 = ||w_n||_V^2 = \int f''(w_n)w_n^2,$$

because $w_n \in \mathcal{V}$. Furthermore, by hypothesis, we have

$$||w_n||_V = \int f'(w_n)\frac{w_n}{||w_n||_V} + \sigma_n.$$

Thus, by f_μ,

$$\begin{aligned} \mu_2 ||w_n||_V^2 &= \mu_2 \int f'(w_n)w_n + \mu_2 \sigma_n ||w_n||_V < \\ &< \int f''(w_n)w_n^2 + \mu_2 \sigma_n ||w_n||_V = \\ &= ||w_n||_V^2 + \mu_2 \sigma_n ||w_n||_V. \end{aligned}$$

So, because $\mu_2 > 1$ we have that

$$0 < (\mu_2 - 1)||w_n||_V < \mu_2 \sigma_n \to 0, \tag{18}$$

that is a contradiction. □

Remark 10. Obviously, by Lemma 8 we have also

$$B := \inf_{||u||_V = 1} T_u > 0,$$

and B does not depend on g.

At last we can give the following characterization of the Nehari manifold.

Proposition 11. *Let $||g||_{L^{\frac{2N}{N+2}}}$, sufficiently small, and let $u \in \mathcal{D}^{1,2}$ with $||u||_V = 1$. Then*

1. *If* $\int gu < 0$, *then there exists an unique* t_u^1 *such that* $t_u^1 u \in \mathcal{N}_g^V$ *and* $t_u^0 < t_u^1$, *where* t_u^0 *is the unique value for which* $t_u^0 \in \mathcal{N}_0^V$.

2. *If* $\int gu = 0$, *then there exists an unique* t_u^1 *such that* $t_u^1 u \in \mathcal{N}_g^V$ *and* $t_u^0 = t_u^1$.

3. *If* $\int gu > 0$, *then there exist two positive numbers* t_u^1 *and* t_u^2 *such that* $t_u^j u \in \mathcal{N}_g^V$ *and* $t_u^2 < T_u < t_u^1 < t_u^0$, *where* T_u *is the unique value for which* $T_u u \in \mathcal{V}$.

4. t_u^1 *and* t_u^2 *depend* C^1 *on* $g \in L^{\frac{2N}{N+2}}$ *and on* $u \in \mathcal{D}^{1,2} \smallsetminus \{0\}$. *Furthermore, fixed* u, *we have* $t_u^1 \to t_u^0$, *when* $||g||_{L^{\frac{2N}{N+2}}} \to 0$.

Proof. 1. If $\varphi_g'(\bar{t}) = 0$, with $\bar{t} \neq 0$, by f_μ, we have that

$$\bar{t}^2 \varphi_g''(\bar{t}) = \bar{t} \int gu + \int [\bar{t} u f'(\bar{t}u) - \bar{t}^2 u^2 f''(\bar{t}u)] < 0, \tag{19}$$

so $\bar{t}$ is a maximum point for φ_g. Furthermore we have that $\varphi_g(0) = 0$, $\varphi_g'(0) > 0$ and $\varphi_g''(0) > 0$.

Using (f_μ) and (f_0), we have

$$\begin{aligned}
\varphi_g(t) &= \frac{t^2}{2}||u||_V^2 - \int_{\mathbb{R}^N} f(tu) - t \int_{\mathbb{R}^N} gu \leq \\
&\leq \frac{t^2}{2}||u||_V^2 - t \int_{\mathbb{R}^N} gu - c_0 t^q \int_{t|u|<1} |u|^q - c_0 t^q \int_{t|u|\geq 1} |u|^p \leq \qquad (20)\\
&\leq \frac{t^2}{2}||u||_V^2 - t \int_{\mathbb{R}^N} gu - c_0 t^p \int_{|u|\geq 1} |u|^p \to -\infty \text{ when } t \to \infty,
\end{aligned}$$

because $p > 2$. This proves that there is exactly one t_u^1 such that $t_u^1 u \in \mathcal{N}_g$; it is easy to see that $t_u^0 < t_u^1$.

2. In this case, we can proof, as in (20) that $\varphi_g(t) \to -\infty$ when $t \to \infty$ and that if $\bar{t} \neq 0$ is a critical point of φ_g then (19) holds. At last, consider that $0 = \varphi_g(0) = \varphi_g'(0) < \varphi_g''(0)$, and so 0 is a local minimum for φ_g, and we can conclude.

3. We have just proved that, for any $u \in \mathcal{D}^{1,2}$, we have an unique maximum point T_u of $\varphi_0'(t)$. So, if we prove that $\int gu < \varphi_0'(T_u)$ we have that there exist two numbers t_u^1 and t_u^2 such that $\varphi_g'(t_u^j) = 0$. Set L as in Lemma 9, and consider that

$$\int gu \leq ||g||_{L^{\frac{2N}{N+2}}} ||u||_{L^{2^*}} \leq C_1 ||g||_{L^{\frac{2N}{N+2}}} ||u||_{\mathcal{D}^{1,2}} \leq C_2 ||g||_{L^{\frac{2N}{N+2}}} ||u||_V. \tag{21}$$

Recalling that $||u||_V = 1$, if $||g||_{L^{\frac{2N}{N+2}}}$ is sufficiently small, that is $C_2||g||_{L^{\frac{2N}{N+2}}} < L$, we have exactly two positive numbers t_u^1 and t_u^2 such that $\varphi_g'(t_u^j) = 0$, and t_u^1 and t_u^2 are respectively the maximum and the minimum point of φ_g

4. For Simplicity we only prove that $t_u^1(g)$ is a C^1 function. The other case is straightforward. Let us define a function $G : \mathbb{R}^+ \times \mathcal{D}^{1,2} \setminus \{0\} \times L^{\frac{2N}{N+2}} \to \mathbb{R}$,

$$G : (t, u, g) \mapsto \frac{d}{dt}\varphi_g^u(t) = t||u||_V^2 - \int f'(tu)u - \int gu.$$

We have that G is a C^1 function. Let $\bar{t}, \bar{u}, \bar{g}$ be such that $G(\bar{t}, \bar{u}, \bar{g}) = 0$. We know that $\frac{\partial}{\partial t}G(\bar{t}, \bar{u}, \bar{g}) = \frac{d^2}{dt^2}\varphi_{\bar{g}}^{\bar{u}}(\bar{t}) < 0$, thus, by the implicit function theorem there is a C^1 function $t(u, g) = t_u^1(g)$ such that $G(t(u, g), u, g) = 0$. We have then the claimed result. $\square$

The Nehari manifold so can be described as:

$$\mathcal{N}_g^V = \mathcal{N}_{g,V}^+ \cup \mathcal{N}_{g,V}^-, \tag{22}$$

where

$$\begin{aligned} \mathcal{N}_g^+ &= \mathcal{N}_{g,V}^+ := & \{u \in \mathcal{N}_g^V \ : \ E_g'(u)u = 0, E_g''(u)u^2 > 0\}; \\ \mathcal{N}_g^- &= \mathcal{N}_{g,V}^- := & \{u \in \mathcal{N}_g^V \ : \ E_g'(u)u = 0, E_g''(u)u^2 < 0\}. \end{aligned}$$

We have also that $E_g^V > 0$ on $\mathcal{N}_g^-$ and $E_g^V < 0$ on $\mathcal{N}_g^+$. Furthermore, because $\mathcal{N}_0^V$ and $\mathcal{V}$ are bounded away from 0, we have also that $\inf_{u \in \mathcal{N}_g^-} ||u|| > 0$. The geometry of $\mathcal{N}_g^V$ is represented in the following picture.



Remark 12. There exists $M > 0$ such that

$$||u||_V \leq M||g||_{\frac{2N}{N+2}} \text{ for any } u \in \mathcal{N}_g^+, \tag{23}$$

indeed, by (f_μ) we have

$$\frac{1}{2}||u||_V^2 < \int f(u) + \int gu \leq \frac{1}{\mu_1}\int f'(u)u + \int gu = \frac{1}{\mu_1}||u||_V^2 + \left(1 - \frac{1}{\mu_1}\right)\int gu,$$

so

$$\left(\frac{1}{2} - \frac{1}{\mu_1}\right)||u||_V^2 < \left(1 - \frac{1}{\mu_1}\right)\int gu.$$

3 The Splitting Lemma

We recall that a sequence $\{u_n\}_n \in \mathcal{D}^{1,2}$ such that $E_g^V(u_n) \to c$, and $\nabla E_g^V(u_n) \to 0$ is a Palais-Smale sequence at level c for E_g^V.

In the same way we say that $\{u_n\}_n \in \mathcal{N}_g^V$ such that $E_g^V(u_n) \to c$, and there exists a sequence $\varepsilon_n \to 0$ s.t. $|\langle \nabla E_g^V(u_n), \varphi \rangle| \leq \varepsilon_n ||\varphi||$, for all $\varphi \in T_{u_n}\mathcal{N}_g^V \cap \mathcal{D}^{1,2}$ is a Palais-Smale sequence at level c for E_g^V restricted to $\mathcal{N}_g^V$.

A functional f satisfies the $(PS)_c$ condition if all the Palais-Smale sequences at level c converge.

Unfortunately the functional E_g^V on $\mathcal{N}_g^V$ does not satisfy the PS condition in all the energy range. In this section by the splitting lemma we get a description of the PS sequences for the functional E_g^V.

Lemma 13. *Let $u_n \in \mathcal{N}_g$ and let $E_g^V(u_n) \to c$. Then $||u_n||_V$ is bounded.*

Proof. We have that

$$||u_n||_V^2 = \int f'(u_n)u_n + \int gu_n \tag{24}$$

because $u_n \in \mathcal{N}_g^V$. Furthermore, for (f_μ) we have

$$\begin{aligned} E_g^V(u_n) &= \frac{1}{2}||u_n||_V^2 - \int f(u_n) - \int gu_n \geq \\ &\geq \frac{1}{2}||u_n||_V^2 - \frac{1}{\mu_1}\int f'(u_n)u_n - \int gu_n = \\ &= \frac{1}{2}||u_n||_V^2 - \frac{1}{\mu_1}||u_n||_V^2 + \frac{1}{\mu_1}\int gu_n - \int gu_n = \\ &= \left(\frac{1}{2} - \frac{1}{\mu_1}\right)||u_n||_V^2 - \left(1 - \frac{1}{\mu_1}\right)\int gu_n = \\ &= ||u_n||_V^2 \left[\left(\frac{1}{2} - \frac{1}{\mu_1}\right) - \left(1 - \frac{1}{\mu_1}\right)\int g\frac{u_n}{||u_n||_V^2}\right]. \end{aligned} \tag{25}$$

If $||u_n||_V \to \infty$ we have that

$$\left| \int g \frac{u_n}{||u_n||_V^2} \right| \leq ||g||_{L^{\frac{2N}{N+2}}} \frac{||u||_{L^{2^*}}}{||u_n||_V^2} \leq C ||g||_{L^{\frac{2N}{N+2}}} \frac{1}{||u_n||_V} \to 0. \tag{26}$$

So we will have

$$C_1 > E_g^V(u_n) \geq C_2 ||u_n||_V^2 \to \infty \tag{27}$$

that is a contradiction. □

Lemma 14. *Let* $\{u_n\}_n \subset \mathcal{N}_g$*, and let* $E_g^V(u_n) \to c$*. Then, up to subsequence* $u_n \rightharpoonup u_0$ *in* $\mathcal{D}^{1,2}$*. Furthermore, setting* $\psi_n = u_n - u_0$ *we have*

1. $||\psi_n||_V^2 = ||u_n||_V^2 - ||u_0||_V^2 + o(1)$;
2. $E_g^V(\psi_n) = E_g^V(u_n) - E_g^V(u_0) + o(1)$.

Proof. By the previous lemma we have that $||u_n||_{\mathcal{D}^{1,2}}$ is bounded. Then $u_n \rightharpoonup u_0$ and we have that

$$||\psi_n||_V^2 = ||u_n||_V^2 - ||u_0||_V^2 + o(1).$$

Furthermore, we have that

$$\int f(\psi_n) = \int f(u_n) - \int f(u_0) + o(1). \tag{28}$$

Indeed, we have the following equation, where $\tau, \theta, \sigma \in (0,1)$

$$\begin{aligned} & \int f(u_n) - \int f(u_0) - \int f(\psi_n) = \\ = & \int_{B_R} f(u_0 + \psi_n) - f(u_0) - \int_{B_R^C} f(u_0) + \\ & + \int_{B_R^C} f(u_0 + \psi_n) - f(\psi_n) - \int_{B_R} f(\psi_n) = \\ = & \int_{B_R} f'(u_0 + \tau\psi_n)\psi_n - \int_{B_R^C} f(u_0) + \int_{B_R^C} f'(\theta u_0 + \psi_n) u_0 - \int_{B_R} f'(\sigma\psi_n)\psi_n. \end{aligned}$$

Using Lemma 3 we have that the terms in B_R^C are arbitrarily small when R is sufficiently large. Furthermore, since $\psi_n \to 0$ in $L^p(\Omega)$ for all $\Omega \subset \mathbb{R}^N$ bounded and for all $p < 2^*$, we get that

$$\int f(u_n) - \int f(u_0) - \int f(\psi_n) \to 0.$$

The proof follows easily. □

Lemma 15. *Suppose that* $\psi_n \rightharpoonup 0$ *in* $\mathcal{D}^{1,2}$. *Then we have*

$$\int V\psi_n^2 \to 0 \tag{29}$$

$$\int g\psi_n \to 0 \tag{30}$$

Proof. Again we use that $\psi_n \to 0$ in $L^p(\Omega)$ for all $\Omega \subset \mathbb{R}^N$ bounded and for all $p < 2^*$. We have that

$$\begin{aligned}\int V\psi_n^2 &= \int\limits_{B_R} V\psi_n^2 + \int\limits_{\mathbb{R}^N \smallsetminus B_R} V\psi_n^2 \leq ||V||_{L^t(B_R)}||\psi_n||^2_{L^{2t'}(B_R)} + \\ &\quad +||V||_{L^{N/2}(\mathbb{R}^N \smallsetminus B_R)}||\psi_n||^2_{L^{2^*}(\mathbb{R}^N \smallsetminus B_R)} \to 0,\end{aligned}$$

and that

$$\begin{aligned}\int g\psi_n &= \int\limits_{B_R} g\psi_n + \int\limits_{\mathbb{R}^N \smallsetminus B_R} g\psi_n \leq ||g||_{L^s(B_R)}||\psi_n||_{L^{s'}(B_R)} + \\ &\quad +||g||_{L^{\frac{2N}{N+2}}(\mathbb{R}^N \smallsetminus B_R)}||\psi_n||_{L^{2^*}(\mathbb{R}^N \smallsetminus B_R)} \to 0.\end{aligned}$$

□

Lemma 16. *Let* $\{u_n\}_n$ *a PS sequence at level* c *for the functional* E_g^V *restricted to the manifold* $\mathcal{N}_g^V$. *Then, up to a subsequence, there exist* k *sequences of points* $\{y_n^j\}_n$, $j = 1, \dots k$, *with* $|y_n^j| \to \infty$, *a solution* u^0 *of the problem* $-\Delta u + Vu = f'(u) + g$, *and* k *solutions* u^j, $j = 1, \dots k$, *of the problem* $-\Delta u = f'(u)$ *such that*

$$u_n(x) = u^0(x) + \sum_{j=1}^k u^j(x - y_n^j) + o(1); \tag{31}$$

$$E_g^V(u_n) = E_g^V(u^0) + \sum_{j=1}^k E_0^0(u^j) + o(1). \tag{32}$$

Proof. Since u_n is a *PS* sequence for the functional E_g^V restricted to the manifold $\mathcal{N}_g^V$, then u_n is a *PS* sequence for the functional E_g^V. By the Lemma 14 we have that u_n converges to u^0 weakly in $\mathcal{D}^{1,2}$ (up to subsequence), so, given $\varphi \in C_0^\infty(\mathbb{R}^N)$,

$$\lim_{n\to\infty} \int \nabla u_n \nabla\varphi + Vu_n\varphi - f'(u_n)\varphi - g\varphi = 0. \tag{33}$$

It is easy to see that

$$\int \nabla u_n \nabla \varphi + V u_n \varphi \to \int \nabla u^0 \nabla \varphi + V u^0 \varphi.$$

Arguing as in Step 1 of [5, Lemma 3.3] we get also that, for some $0 < \theta < 1$,

$$\int [f'(u_n) - f'(u^0)]\varphi = \int_{\text{supp}\varphi} f''(\theta u_n + (1-\theta)u^0)(u_n - u^0)\varphi \to 0, \tag{34}$$

as $n \to 0$, because $u_n - u^0 \to 0$ in $L^p(\Omega)$, with Ω bounded and $p < 2^*$. So we have proved that u^0 solves $-\Delta u + Vu = f'(u) + g$.

Now we set

$$\psi_n(x) = u_n(x) - u^0(x).$$

Then $\psi_n \rightharpoonup 0$ weakly in $\mathcal{D}^{1,2}$. If $\psi_n \nrightarrow 0$ strongly in $\mathcal{D}^{1,2}$, for Step 3 of [5, Lemma 3.3] we have that there exists a sequence $\{y_n\} \subset \mathbb{R}^N$ with $|y_n| \to \infty$ such that $\psi_n(x + y_n) \to u^1$ in $\mathcal{D}^{1,2}$, and $u^1 \neq 0$.

Because u^0 is a weak solution of $(\mathscr{P})$ and u_n is a PS sequence for E_g^V we have that, for any $\varphi \in C_0^\infty(\mathbb{R}^N)$,

$$\int \nabla u_n \nabla \varphi + V u_n \varphi - f'(u_n)\varphi - g\varphi \to 0;$$

$$\int \nabla u^0 \nabla \varphi + V u^0 \varphi - f'(u^0)\varphi - g\varphi = 0.$$

So

$$\int \nabla \psi_n \nabla \varphi + V \psi_n \varphi - f'(u_n) - f'(u^0)\varphi \to 0. \tag{35}$$

Using (34) we have that ψ_n is a PS sequence for the functional E_0^V. Thus, for any $\varphi \in C_0^\infty(\mathbb{R}^N)$ we have

$$\begin{aligned}
&\int \nabla \psi_n(x + y_n) \nabla \varphi(x) - f'(\psi_n(x + y_n))\varphi(x)dx = \\
&\int \nabla \psi_n(x) \nabla \varphi(x - y_n) - f'(\psi_n(x))\varphi(x - y_n)dx = \\
&\int [f'(u_n) - f'(u^0) - f'(\psi_n)]\varphi(x - y_n) - \int V(x)\psi_n(x)\varphi(x - y_n) + o(1).
\end{aligned}$$

Using the same argument of Lemma 15 we can prove that

$$\int V(x)\psi_n(x)v(x - y_n) \leq C\varepsilon_n ||\varphi||_{\mathcal{D}^{1,2}}, \text{ with } \varepsilon_n \to 0;$$

furthermore we have

$$\begin{aligned}
&\int [f'(u_n) - f'(u^0) - f'(\psi_n)]\varphi(x-y_n) = \\
= &\int_{B_R} [f'(u_0+\psi_n) - f'(u^0)]\varphi(x-y_n) + \\
&+ \int_{B_R^C} [f'(u_0+\psi_n) - f'(\psi_n)]\varphi(x-y_n) - \\
&- \int_{B_R^C} f'(u^0)\varphi(x-y_n) + \int_{B_R} f'(\psi_n)\varphi(x-y_n) \le \\
\le &\ ||[f''(u^0+\theta\psi_n) - f''(\theta\psi_n)]\varphi(\cdot - y_n)||_{L^{p'}(\mathbb{R}^N)}||\psi_n||_{L^p(B_R)} + \\
&+||[f''(\psi_n+\theta u^0) - f''(\theta u^0)]\varphi(\cdot - y_n)||_{L^{p'}\cap L^{q'}}||u^0||_{L^p+L^q(B_R^C)},
\end{aligned}$$

where $0 < \theta < 1$. Because $||u^0||_{L^p+L^q(B_R^C)} \to 0$ for $R \to \infty$, and given R $||\psi_n||_{L^p(B_R)} \to 0$ as $n \to 0$, we have that

$$\int \nabla\psi_n(x+y_n)\nabla\varphi(x) - f'(\psi_n(x+y_n))\varphi(x)dx \to 0.$$

At last, it is easy to see that

$$\int \nabla\psi_n(x+y_n)\nabla v(x) - f'(\psi_n(x+y_n))v(x)dx \to \int \nabla u^1\nabla v(x) - f'(u^1)v(x)dx,$$

so we have also proved that u^1 solves the problem $-\Delta u = f(u)$.

Set $\psi_n^2 = \psi(x+y_n) - u^1$, we have that $\psi_n^2 \rightharpoonup 0$, thus

$$\begin{aligned}
E_g^V(u_n) - E_g^V(u^0) &= E_g^V(\psi_n(x)) + o(1) = E_0^0(\psi_n(x)) + o(1) = \\
&= E_0^0(\psi_n(x+y_n)) + o(1) = E_0^0(u^1) + E_0^0(\psi_n^2) + o(1),
\end{aligned}$$

by Lemma 14. So,

$$E_g^V(u_n) = E_g^V(u^0) + E_0^0(u^1) + E_0^0(\psi_n^2) + o(1). \tag{36}$$

Now, if $\psi_n^2 \to 0$ strongly in $\mathcal{D}^{1,2}$, we have the claim, otherwise we can proceed by induction and conclude the proof in a finite number of steps. □

4 Main Results

We set

$$m_g = \inf_{u\in\mathcal{N}_g^V} E_g^V(u) \text{ and } m_{1,g} = \inf_{u\in\mathcal{N}_g^-} E_g^V(u).$$

We show that there exist a solution with critical value m_g and another solution with critical value $m_{1,g}$.

We set also

$$m_0 = \inf_{u\in\mathcal{N}_0^0} E_0^0(u) \tag{37}$$

and we recall that there exists a positive radially symmetric function $\omega \in \mathcal{N}_0^0$ such that

$$E_0^0(\omega) = m_0 > 0. \tag{38}$$

Finally, we set

$$m_V = \inf_{u\in\mathcal{N}_0^V} E_0^V(u) \tag{39}$$

We know, by [5], that for any $V \leq 0$ and $V < 0$ on a set of positive measure there exists a function $\bar{u} \in \mathcal{N}_0^V$ such that

$$E_0^V(\bar{u}) = m_V \tag{40}$$

and

$$0 < m_V < m_0. \tag{41}$$

We prove the following results.

Theorem 17. *There exist a $u_g \in \mathcal{N}_g^+$ such that $E_g^V(u_g) = m_g$. Furthermore, when $||g||_{L^{\frac{2N}{N+2}}}$ is small, u_g is unique.*

Proof. By definition of $\mathcal{N}_g^+$ we have that $m_g = \inf\limits_{u\in\mathcal{N}_g^+} E_g^V(u)$, and that $m_g < 0$. At first we prove that $m_g > -\infty$. By contradiction, suppose that there exist a sequence $t_n > 0$ and a sequence $\{v_n\}_n \subset \mathcal{D}^{1,2}$ with $||v_n||_V = 1$ and $t_n v_n \in \mathcal{N}_g^+$ such that

$$E_g^V(t_n v_n) = \frac{t_n^2}{2} - \int f(t_n v_n) - t_n \int g v_n \to -\infty. \tag{42}$$

We have also that $t_n^2 - \displaystyle\int f'(t_n v_n) t_n v_n - t_n \int g v_n = 0$. So, if t_n is bounded, we have

$$\begin{aligned} E_g^V(t_n v_n) &= -\frac{t_n^2}{2} + \int f'(t_n v_n) t_n v_n - \int f(t_n v_n) \geq \\ &\geq -\frac{t_n^2}{2} + \left(1 - \frac{1}{\mu_1}\right) \int f'(t_n v_n) t_n v_n \end{aligned}$$

that is bounded by Lemma 3. Thus we have that, up to subsequence, $t_n \to +\infty$. Finally, arguing as in (25) we have that

$$E_g^V(t_n v_n) \geq \left(\frac{1}{2} - \frac{1}{\mu_1}\right) t_n^2 - \left(1 - \frac{1}{\mu_1}\right) t_n \int g v_n \to +\infty, \tag{43}$$

that is a contradiction.

Now, let u_n a minimizing sequence. For the Ekeland variational principle, we can suppose u_n be a PS sequence. For the splitting lemma there exists a $u_g \in \mathcal{N}_g^V$ and k functions u^j, $1 \leq j \leq k$ such that

$$E_g^V(u_n) \to E_g^V(u_g) + \sum_{j=1}^{k} E_0^0(u^j) = m_g < 0. \tag{44}$$

We know that $E_0^0(u^j) \geq m_0 > 0$ for all j. So, if $k > 0$ we will have $E_g^V(u_n) \to m_g + \delta$ for some $\delta > 0$ and this is a contradiction.

So, we have

$$u_n \to u_g \text{ in } \mathcal{D}^{1,2}. \tag{45}$$

Furthermore, we have $E_g^V(u_g) = m_g < 0$, so $u_g \in \mathcal{N}_g^+$, and this concludes the proof of the existence.

To prove uniqueness, we argue by contradiction. If u_1, u_2 are minimizers of E_g^V on $\mathcal{N}_g^+$, both u_1 and u_2 solve ($\mathscr{P}$), so we have

$$||u_1 - u_2||_V^2 = \int (f'(u_1) - f'(u_2))(u_1 - u_2) = \int f''(\theta u_1 + (1-\theta)u_2)(u_1 - u_2)^2$$

with $0 < \theta < 1$. So

$$||u_1-u_2||_{L^{2^*}}^2 \leq C||u_1-u_2||_V^2 \leq C||u_1-u_2||_{L^{2^*}}^2 ||f''(\theta u_1+(1-\theta)u_2)||_{L^{\frac{2^*}{2^*-2}}}. \tag{46}$$

By Remark 12, we have that, if $g \to 0$ in $L^{\frac{2N}{N+2}}$, then both u_1 and u_2 are small in $L^p + L^q$, so we have that $f''(\theta u_1 + (1-\theta)u_2) \to 0$ in $L^{p/p-2} \cap L^{q/q-2}$ by Lemma 5, and, by interpolation,

$$||f''(\theta u_1 + (1-\theta)u_2)||_{L^{\frac{2^*}{2^*-2}}} \to 0,$$

that is a contradiction. □

Proposition 18. *Suppose that $g \geq 0$. Then there exists an $u_g \geq 0$ in $\mathcal{N}_g^+$ such that $E_g^V(u_g) = m_g$.*

Proof. Take u_g as in Theorem 17. Because $u_g \in \mathcal{N}_g^+$ we have that $\int g u_g > 0$. If u_g changes sign, or u_g negative, we have that

$$0 < \int g u_g \leq \int g|u_g|. \tag{47}$$

So, reminding that f is even we have

$$\begin{aligned} E_g^V(|u_g|) &= \frac{1}{2}||u_g||_V^2 - \int f(|u_g|) - \int g|u_g| \leq \\ &\leq \frac{1}{2}||u_g||_V^2 - \int f(u_g) - \int g u_g = E_g^V(u_g). \end{aligned}$$

We know that there exists a τ such that $\tau|u_g| \in \mathcal{N}_g^+$. Furthermore we know, by the study of $\varphi_g^{|u_g|}$ that τ is a local minimizer of $\varphi_g^{|u_g|}$, in fact, $\varphi_g^{|u_g|}(\tau) \leq \varphi_g^{|u_g|}(t)$ for all $t \in [0, \tau]$. We have

$$\begin{aligned} \frac{d}{dt}\varphi_g^{|u_g|}(1) &= \frac{d}{dt}E_g^V(t|u_g|)_{|t=1} = ||u_g||_V^2 - \int f'(|u_g|)|u_g| - \int g|u_g| \leq \\ &\leq ||u_g||_V^2 - \int f'(u_g)u_g - \int g u_g = \frac{d}{dt}E_g^V(tu_g)_{|t=1} = 0, \end{aligned}$$

and

$$\begin{aligned} \frac{d^2}{dt^2}\varphi_g^{|u_g|}(1) &= \frac{d^2}{dt^2}E_g^V(t|u_g|)_{|t=1} = ||u_g||_V^2 - \int f''(|u_g|)|u_g|^2 = \\ &= ||u_g||_V^2 - \int f''(u_g)u_g^2 = \frac{d^2}{dt^2}E_g^V(tu_g)_{|t=1} > 0. \end{aligned}$$

Thus $\tau \geq 1$ and

$$E_g^V(\tau|u_g|) \leq E_g^V(|u_g|) \leq E_g^V(u_g) = m_g, \tag{48}$$

that concludes the proof. $\square$

We want to prove that, under suitable hypothesis on g, f and V, there exists another solution of $\mathscr{P}$, by minimizing the functional E_g^V on $\mathcal{N}_g^-$. In order to prove that a minimizing sequence converges we will show that, for g small,

$$m_{1,g} := \inf_{u \in \mathcal{N}_g^-} E_g^V(u) < m_g + m_0; \tag{49}$$

Lemma 19. *Suppose that* $V \leq 0$ *and* $V < 0$ *on a set of positive measure. If* $||g||_{L^{\frac{2N}{N-2}}}$ *sufficiently small, then there exist a* $\delta > 0$ *such that*

$$m_{1,g} := \inf_{u \in \mathcal{N}_g^-} E_g^V(u) < m_0 - \delta. \tag{50}$$

Moreover,

$$\limsup_{||g||_{L^{\frac{2N}{N-2}}} \to 0} m_{1,g} \leq m_V \tag{51}$$

Proof. By [5, Lemma 4.4(a)] and [5, Theorem 1.1] we know that there exists a $\bar{u} \in \mathcal{N}_0^V$ such that

$$E_0^V(\bar{u}) = \inf_{u \in \mathcal{N}_0^V} E_0^V(u) = m_V < m_0.$$

We set $v = \frac{\bar{u}}{||\bar{u}||_V}$, so $\bar{u} = t_0^v v$. We know that there exists $t_1^v = t_1^v(g)$ such that $t_1^v v \in \mathcal{N}_g^-$ by Proposition 11. Furthermore, by Proposition 11 we have that $t_1^v \to t_0^v$ when $||g||_{L^{\frac{2N}{N-2}}} \to 0$, and so

$$m_{1,g} \leq E_g^V(t_1^v v) \to E_0^V(\bar{u}) = m_V < m_0 \text{ for } ||g||_{L^{\frac{2N}{N+2}}} \to 0, \tag{52}$$

that concludes the proof. □

Theorem 20. *For* $||g||_{L^{\frac{2N}{N-2}}} \to 0$ *there exist* $u_{1,g} \in \mathcal{N}_g^-$ *a solution of* ($\mathscr{P}$). *Furthermore, if* $g \geq 0$ *the solution* $u_{1,g}$ *can be chosen positive.*

Proof. By the splitting lemma, to obtain the result it is enough to show that $m_{1,g} < m_g + m_0$. In the previous lemma, we have proved that there exists a $\delta > 0$ such that $m_{1,g} < m_0 - \delta$ for $||g||_{L^{\frac{2N}{N-2}}}$ sufficiently small. By Remark 11 we have also that $m_g \to 0$ when $g \to 0$ in $L^{\frac{2N}{N+2}}$. So there exists $u_{1,g} \in \mathcal{N}_g^V$ a solution of ($\mathscr{P}$). Moreover $E_g^V(u_{1,g})$ is positive, so $u_{1,g} \in \mathcal{N}_V^-$.

To prove the last claim, consider that $E_g^V(|u_{1,g}|) \leq E_g^V(u_{1,g})$. Also, there exists a $\bar{t}$ such that $\bar{t}|u_{1,g}| \in \mathcal{N}_g^-$. Then we have

$$m_{1,g} = E_g^V(u_{1,g}) = \max_t E_g^V(tu_{1,g}) \geq E_g^V(\bar{t}u_{1,g}) \geq E_g^V(\bar{t}|u_{1,g}|). \tag{53}$$

So if $u_{1,g}$ is a solution, also $\bar{t}|u_{1,g}| \in \mathcal{N}_g^-$ is a solution of ($\mathscr{P}$). □

Proposition 21. *If* $||g||_{L^{p'} \cap L^{q'}} \to 0$, *then* $m_{1,g} \to m_V$.

Proof. We take a sequence of $g_n \to 0$ in $L^{p'} \cap L^{q'}$. We know that for any g_n there exists u_{1,g_n} such that $E_{g_n}^V(u_{1,g_n}) = m_{1,g_n}$. For simplicity we call $u_n = u_{1,g_n}$. Also, we set $v_n = \frac{u_n}{||u_n||_{L^p + L^q}}$, and $u_n = t_n v_n$. We have

$$E_{g_n}^V(u_n) = t_n \left[\frac{1}{2} \int f'(t_n v_n) v_n - \int \frac{f(t_n v_n)}{t_n} - \frac{1}{2} \int g_n v_n, \right] \tag{54}$$

and we have that there exist a $\delta > 0$ such that $0 \leq E_{g_n}^V(u_n) \leq m_v + \delta$. Now, suppose, by contradiction, that $t_n \to \infty$. Then,

$$\frac{1}{2} \int f'(t_n v_n) v_n - \int \frac{f(t_n v_n)}{t_n} - \frac{1}{2} \int g_n v_n \to 0, \tag{55}$$

and so
$$\frac{1}{2}\int f'(t_n v_n)v_n - \int \frac{f(t_n v_n)}{t_n} \to 0. \tag{56}$$
By (f_μ), we have that
$$\begin{aligned}\int f'(t_n v_n)v_n - 2\int \frac{f(t_n v_n)}{t_n} &= \int f'(t_n v_n)v_n - \mu_1 \int \frac{f(t_n v_n)}{t_n} + \\ &\quad +(\mu_1 - 2)\int \frac{f(t_n v_n)}{t_n} \geq \\ &\geq (\mu_1 - 2)\int \frac{f(t_n v_n)}{t_n}.\end{aligned}$$

So $\displaystyle\int \frac{f(t_n v_n)}{t_n} \to 0$. Now the hypothesis on f
$$0 \leq c_0 t_n^{p-1}\left[\int\limits_{|v_n>1|} |v_n|^p + \int\limits_{|v_n<1|} |v_n|^q\right] \leq \int \frac{f(t_n v_n)}{t_n} \to 0, \tag{57}$$
so we have that both $\displaystyle\int\limits_{|v_n|>1} |v_n|^p$ and $\displaystyle\int\limits_{|v_n|<1} |v_n|^q$ vanish when $n \to \infty$, and so
$$1 = ||v_n||_{L^p+L^q} \leq \max\left\{\int\limits_{|v_n|>1} |v_n|^p, \int\limits_{|v_n|<1} |v_n|^q\right\} \to 0 \tag{58}$$
that is a contradiction. Furthermore, by Proposition 11, we have t_n bounded away from 0. So, we have that there exists two positive constants c_1 and c_2 such that
$$0 < c_1 \leq t_n = ||u_n||_{L^p+L^q} \leq c_2 < \infty. \tag{59}$$

Now, let τ_n such that $\tau_n u_n \in \mathcal{N}_0^V$. We can show that $\tau_n \to 1$ when $n \to \infty$. The main idea is that
$$\frac{d}{dt}\varphi_{g_n}^{u_n}(\tau_n) - \frac{d}{dt}\varphi_0^{u_n}(\tau_n) = \int g_n u_n \to 0 \tag{60}$$
because $||u_n||_{L^p+L^q}$ is bounded and $g_n \to 0$ in $L^{p'} \cap L^{q'}$. The details are omitted for the sake of simplicity.

Now we have that
$$E_0^V(\tau_n u_n) - E_0^V(u_n) \to 0. \tag{61}$$

We have that $E^V_{g_n}(u_n)$ is bounded, so, up to subsequences, there exists a d such that $E^V_{g_n}(u_n) \to d$ when $n \to \infty$, and, because u_n is bounded in $L^p + L^q$, also $E^V_0(u_n) \to d$, and, by (61), $E^V_0(\tau_n u_n) \to d$.

So, $d \geq m_V$. By Lemma 19 we know also that $d \leq m_V$ so we get the claim. □

Proof of Theorem 1. By theorems 17 and 20, we have that there exists a $u_g \in \mathcal{N}^+_g$ and $u_{1,g} \in \mathcal{N}^-_g$ that solve $(\mathscr{P})$. Furthermore, by Theorem 20 and Proposition 18 the solution can be chosen nonnegative. At least, by Remark 12 we have that $u_g \to 0$ in $\mathcal{D}^{1,2}$ and by Proposition 21 that $m_{1,g} \to m_V$ when $g \to 0$. □

A The Hypothesis on f

We want to prove that there exists a function that satisfies all the conditions required in the introduction.

We take the function

$$f(s) = \frac{|s|^q}{1 + |s|^{q-p}}. \tag{62}$$

This function is even, and it satisfies (f_0).

We have that, for $s > 0$

$$f'(s) = \frac{qs^{q-1} + ps^{2q-p-1}}{(1 + s^{q-p})^2},$$

$$f''(s) = \frac{s^{q-2}}{(1 + s^{q-p})^2} \left\{ q(q-1) + p(2q-p-1)s^{q-p} - \frac{2(q-p)(q+ps^{q-p})s^{q-p}}{1 + s^{q-p}} \right\}.$$

It's easy to see that f satisfies (f_2) and the first part of (f_μ).

We set $\mu_2 = 1+\varepsilon > 1$; then the inequality $(1+\varepsilon)f'(s)s < f''(s)s^2$ becomes

$$(q^2 - 2q - \varepsilon q) + p(2q - p - 2 - \varepsilon)\gamma - \frac{2(q-p)(q+p\gamma)\gamma}{1+\gamma} > 0,$$

where $\gamma = s^{q-p}$. So, we have to prove that

$$q(q-2-\varepsilon) + [p(2q-p-2-\varepsilon) + q(2p-q-2-\varepsilon)]\gamma + p(p-2+\varepsilon)\gamma^2 > 0.$$

Obviously we can choose ε such that $q(q-2-\varepsilon) > 0$ and $p(p-2+\varepsilon) > 0$. Furthermore, we choose $q - p$ sufficiently small such that also $2q - p - 2 - \varepsilon$ and $2p - q - 2 - \varepsilon$ are positive, so the second part of (f_μ) is proved.

At last we prove (f_3) and that $f'''(s)s^3 > 0$. We have that, for $s > 0$,

$$\begin{aligned} f'''(s) &= \frac{6(p-q)^3 s^{4q-3p-3}}{(1+s^{q-p})^4} - \frac{6(1+p-2q)(p-q)^2 s^{3q-2p-3}}{(1+s^{q-p})^3} + \\ &+ \frac{(2p+3p^2+p^3-2q-12pq-6p^2q+9q^2+12pq^2-7q^3)s^{2q-p-3}}{(1+s^{q-p})^2} + \\ &+ \frac{q(2-3q+q^2)s^{q-3}}{1+s^{q-p}}. \end{aligned}$$

We obtain that

$$f'''(s)s^3 = \frac{As^q}{1+s^{q-p}} + \frac{Bs^{2q-p}}{(1+s^{q-p})^2} + \frac{Cs^{3q-2p}}{(1+s^{q-p})^3} + \frac{Ds^{4q-3p}}{(1+s^{q-p})^4},$$

were

$$\begin{array}{ll} A = q(q-2)(q-1); & B = (p-q)(2+3p+p^2-9q-5pq+7q^2); \\ C = 6(p-q)^2(2q-p-1); & D = 6(p-q)^3. \end{array}$$

We can choose $q-p$ sufficiently small, in order to have $B, C, D << A$. Now, set as above $\gamma = s^{q-p}$, we have

$$f'''(s)s^3 = \frac{s^q\left[A+(3A+B)\gamma+(3A+2B+C)\gamma^2+(A+B+C+D)\gamma^3\right]}{(1+s^{q-p})^4}$$

that is positive for all $s > 0$. So (f_μ) is completely proved.

Furthermore, we have that

$$\lim_{s\to 0^+} \frac{f'''(s)}{s^{q-3}} = A = q(q-1)(q-2) > 0, \tag{63}$$

and

$$\lim_{s\to +\infty} \frac{f'''(s)}{s^{p-3}} = A+B+C+D = p(p-1)(p-2) > 0. \tag{64}$$

So, there exists a $c_3 > 0$ such that

$$\begin{cases} |f'''(s)| \le c_3|s|^{p-3} & \text{for } |s| \ge 1; \\ |f'''(s)| \le c_3|s|^{q-3} & \text{for } |s| \le 1. \end{cases} \tag{65}$$

Now, let $\Gamma = \{x \in \mathbb{R}^N \ : \ |u(x)| > 1\}$ and $\Delta = \mathbb{R}^N \smallsetminus \Gamma$ We have that

$$\begin{aligned} \int f'''(u)u^3 &\le \int_\Gamma f'''(u)u^3 + \int_\Delta f'''(u)u^3 \le c_3\int_\Gamma |u|^p + c_3\int_\Delta |u|^q \le \\ &\le C_1 + C_2||u||_{L^p+L^q} \le C_3 + C_4||u||_{\mathcal{D}^{1,2}} < \infty, \end{aligned}$$

and this proves (f_3).

References

[1] Antonio Azzollini and Alessio Pomponio, *Compactness result and applications to some "zero mass" elliptic problems*, ArXiv preprint.

[2] Marino Badiale and Sergio Rolando, *Elliptic problems with singular potential and double-power nonlinearity*, Mediterr. J. Math. **2** (2005), no. 4, 417–436.

[3] Soohyun Bae and Wei-Ming Ni, *Existence and infinite multiplicity for an inhomogeneous semilinear elliptic equation on* $\mathbf{R}^n$, Math. Ann. **320** (2001), no. 1, 191–210.

[4] Vieri Benci and Donato Fortunato, *Towards a unified field theory for classical electrodynamics*, Arch. Ration. Mech. Anal. **173** (2004), no. 3, 379–414.

[5] Vieri Benci, Carlo R. Grisanti, and Anna Maria Micheletti, *Existence and non existence of the ground state solution for the nonlinear schroedinger equations with* $V(\infty) = 0$, Topol. Methods Nonlinear Anal. **26** (2005), 203–219.

[6] Vieri Benci and Anna Maria Micheletti, *Solutions in exterior domains of null mass nonlinear field equations*, Advanced nonlinear studies **6** (2006), no. 2, 171–198.

[7] Henry Berestycki and Pierre-Louis Lions, *Nonlinear scalar field equations. I. Existence of a ground state*, Arch. Rational Mech. Anal. **82** (1983), no. 4, 313–345.

[8] Henry Berestycki and Pierre-Louis Lions, *Nonlinear scalar field equations. II. Existence of infinitely many solutions*, Arch. Rational Mech. Anal. **82** (1983), no. 4, 347–375.

[9] Jöran Bergh and Jörgen Löfström, *Interpolation spaces. An introduction*, Springer-Verlag, Berlin, 1976, Grundlehren der Mathematischen Wissenschaften, No. 223.

[10] Guy Bernard, *An inhomogeneous semilinear equation in entire space*, J. Differential Equations **125** (1996), no. 1, 184–214.

[11] Haim Brezis and Louis Nirenberg, *A minimization problem with critical exponent and non zero data*, Symmetry in Nature, Scuola Normale Superiore, Pisa, 1989, pp. 129–140.

[12] Marco Ghimenti and Anna Maria Micheletti, *Existence of minimal nodal solutions for the nonlinear Schrödinger equations with* $V(\infty) = 0$, Adv. Differential Equations **11** (2006), no. 12, 1375–1396.

[13] Zeev Nehari, *On a class of nonlinear second-order differential equations*, Trans. Amer. Math. Soc. **95** (1960), 101–123.

[14] Lorenzo Pisani, *Remark on the sum of lebesgue spaces*, Preprint of the University of Bari.

[15] Michael Struwe, *A global compactness result for elliptic boundary value problems involving limiting nonlinearities*, Math. Z. **187** (1984), no. 4, 511–517.

[16] G. Tarantello, *On nonhomogeneous elliptic equations involving critical Sobolev exponent*, Ann. Inst. H. Poincaré Anal. Non Linéaire **9** (1992), no. 3, 281–304.

[17] Huan-Song Zhou, *Solutions for a quasilinear elliptic equation with critical Sobolev exponent and perturbations on* $\mathbf{R}^N$, Differential Integral Equations **13** (2000), no. 4-6, 595–612.

[18] Xi Ping Zhu, *A perturbation result on positive entire solutions of a semilinear elliptic equation*, J. Differential Equations **92** (1991), no. 2, 163–178.